



TITLE:

Gallium(III)-catalysed bromocyanation of alkynes: regio- and stereoselective synthesis of  $\beta$ -bromo- $\alpha,\beta$ -unsaturated nitriles.

AUTHOR(S):

Murai, Masahito; Hatano, Ryo; Kitabata, Sachie; Ohe, Kouichi

---

CITATION:

Murai, Masahito ...[et al]. Gallium(III)-catalysed bromocyanation of alkynes: regio- and stereoselective synthesis of  $\beta$ -bromo- $\alpha,\beta$ -unsaturated nitriles.. Chemical communications 2011, 47(8): 2375-2377

ISSUE DATE:

2011-02

URL:

<http://hdl.handle.net/2433/156438>

RIGHT:

© Royal Society of Chemistry 2011.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

# Gallium (III)-catalysed Bromocyanation of Alkynes: Regio- and Stereoselective Synthesis of $\beta$ -Bromo- $\alpha,\beta$ -unsaturated Nitriles

Masahito Murai, Ryo Hatano, Sachie Kitabata, and Kouichi Ohe\*


Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

First published on the web Xth XXXXXXXXXX 20XX

**Treatment of arylacetylenes and cyanogen bromide in  $\text{ClCH}_2\text{CH}_2\text{Cl}$  with a catalytic amount of  $\text{GaCl}_3$  afforded (Z)- $\beta$ -bromoacrylonitriles with high regio- and stereoselectivity.**

10

The catalytic addition reactions of X-Y-type substrates to carbon-carbon multiple bonds are of continued interest due to the facile access to 1,2-difunctional units from simple alkenes or alkynes with ideal atom efficiency.<sup>1</sup> Among these transformations, addition reactions of X-CN to alkynes simultaneously form vinyl-X and vinyl-carbon bonds, both of which can be used to construct complex structures. Several notable examples of palladium- or nickel-catalysed regio- and stereoselective addition reactions to alkynes with several X-CN groups have been reported, such as X=H (hydrocyanation),<sup>2</sup> X=C (carbocyanation),<sup>3</sup> X=Si (cyanosilylation),<sup>4</sup> X=Ge (cyanogermylation),<sup>5</sup> X=Sn (cyanostannylation),<sup>6</sup> X=B (cyanoboration),<sup>7</sup> X=S (cyanothiolation).<sup>8</sup> However, much less attention has been paid to catalytic regio- and stereoselective halocyanation of alkynes<sup>9</sup> or alkenes<sup>10</sup> using cyanogen halides. Herein, we report on gallium-catalysed bromocyanation of alkynes with cyanogen bromide, providing an efficient route to (Z)- $\beta$ -bromoacrylonitriles in a high regio- and stereoselective fashion (Scheme 1). Taking advantage of (Z)- $\beta$ -bromoacrylonitriles, we can establish efficient routes to a wide range of  $\alpha,\beta$ -unsaturated nitriles,<sup>11</sup> which are of synthetic value.



**Scheme 1** Catalytic Addition Reactions of X-CN to Alkynes.

When we examined the reaction of cyanogen bromide<sup>12</sup> and phenylacetylene using palladium or nickel/phosphine complexes, which are effective catalysts in addition reactions of X-CN to alkynes (vide supra), no adducts were generated. Next, Lewis acids were screened for bromocyanation of alkynes, because there is a precedent for the haloacylation of alkynes in analogous reactions.<sup>13</sup> Representative results of the reaction of cyanogen

bromide with phenylacetylene are shown in Table 1. We found that phenylacetylene underwent bromocyanation in the presence of  $\text{AlCl}_3$  (10 mol%) in 1,2-dichloroethane at 80 °C to give  $\beta$ -bromocinnamionitrile **1a** in 42% yield as a mixture of Z- and E-isomers (Z:E = 89:11) (Table 1, entry 1). Interestingly, the use of  $\text{GaCl}_3$  (10 mol%) instead of  $\text{AlCl}_3$  led to **1a** in a high yield and stereoselectivity (81% chemical yield, Z:E = 92:8) (entry 2).<sup>14</sup> This is in sharp contrast with the non-catalysed bromocyanation of ynamines, which gave a low stereoselectivity of the adducts (Z:E = 50:50, ~ 60:40).<sup>9a</sup> The reaction using  $\text{GaCl}_3$  at 70 °C led to a lower yield of **1a**, but with similar stereoselectivity (entry 3). 1,2-Dichloroethane was the most suitable solvent for bromocyanation, while other solvents, e.g.,  $\text{CHCl}_3$ , toluene, heptane, and 2-methyltetrahydrofuran gave a lower yield of the adducts (entries 4-6), or no adducts (entry 7).<sup>15</sup> Using  $\text{GaBr}_3$  as a catalyst afforded almost the same result as  $\text{GaCl}_3$  (entry 8). Other Lewis acid catalysts, such as  $\text{InCl}_3$  and  $\text{InBr}_3$ , showed marginal catalytic activity and gave lower yields of **1a** (entries 9 and 10), while  $\text{FeBr}_3$ ,  $\text{CuBr}_2$ , and  $\text{ZnBr}_2$  exhibited no catalytic activity for bromocyanation (entries 11-13). It should be noted that

**Table 1.** Lewis Acid-catalysed Bromocyanation of Phenylacetylene Using  $\text{BrCN}^a$

$\text{Ph—}\equiv + \text{BrCN} \xrightarrow[\text{solvent, 80 }^\circ\text{C, 12 h}]{\text{catalyst}} \text{Ph—C(Br)=C(R)CN}$ <p style="text-align: center;"><b>1a</b></p>				
entry	catalyst	solvent	yield <sup>b</sup>	Z : E <sup>c</sup>
1	$\text{AlCl}_3$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	42%	89 : 11
2	$\text{GaCl}_3$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	81% (72%)	92 : 8
3 <sup>d</sup>	$\text{GaCl}_3$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	41%	91 : 9
4	$\text{GaCl}_3$	$\text{CHCl}_3$	62%	90 : 10
5	$\text{GaCl}_3$	toluene	61%	90 : 10
6	$\text{GaCl}_3$	heptane	25%	91 : 9
7	$\text{GaCl}_3$	2-MeTHF	0%	—
8	$\text{GaBr}_3$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	75%	96 : 4
9	$\text{InCl}_3$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	15%	93 : 7
10	$\text{InBr}_3$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	32%	95 : 5
11	$\text{FeBr}_3$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	0%	—
12	$\text{CuBr}_2$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	0%	—
13	$\text{ZnBr}_2$	$\text{ClCH}_2\text{CH}_2\text{Cl}$	0%	—

<sup>a</sup> Reaction conditions: Phenylacetylene (0.48 mmol) and  $\text{BrCN}$  (0.40 mmol) in solvent (1.6 mL) were heated in the presence of catalyst (10 mol%). <sup>b</sup> NMR yield (anisole as an internal standard). Isolated yield in parentheses. <sup>c</sup> Determined by NMR. <sup>d</sup> At 70 °C.

Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, Katsura, Nishikyo-ku, Kyoto 615-8510, Japan.

E-mail: ohe@scl.kyoto-u.ac.jp


Tel: (+81) 75-383-2495; Fax: (+81) 75-383-2499

† Electronic Supplementary Information (ESI) is available: Experimental details and analytical data are provided. See DOI: XXXXXXXX

no chlorocyanation adducts were obtained even when Lewis acid catalysts bearing chloride ligands were used (entries 1-7 and 9).

With the optimized reaction conditions established (10 mol% GaCl<sub>3</sub> in 1,2-dichloroethane at 80 °C), we then examined the substrate scope of alkynes (Table 2). Arylacetylenes having a range of aromatic rings underwent bromocyanation of the alkyne moieties to give the corresponding (*Z*)-adducts, **1b-h**<sup>16</sup> in good yields with high regio- and stereoselectivity (entries 1-7), while 1-octyne and 1-(trimethylsilyl)acetylene gave no adducts. Reactions with internal aliphatic or alicyclic alkynes, such as 4-octyne and cyclooctyne, gave complex mixtures, while internal alkynes substituted by a phenyl ring produced bromocyanation adducts **1i**<sup>17</sup> and **1j**, having a cyano group at the β position to the phenyl group in good yields with high regio- and stereoselectivities (entries 8 and 9). Although the reaction of diphenylacetylene was sluggish, and required an elevated temperature (100 °C), the corresponding bromocyanation adduct **1k** was obtained in a 56% yield with excellent stereoselectivity (entry 10).

**Table 2.** GaCl<sub>3</sub>-catalysed Bromocyanation of Alkynes Using BrCN<sup>a</sup>




entry	R <sup>1</sup>	R <sup>2</sup>	product	isolated yield	Z : E <sup>b</sup>
1	4-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	H	<b>1b</b>	70%	95 : 5
2	2-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	H	<b>1c</b>	61%	98 : 2
3	2-naph	H	<b>1d</b>	55%	95 : 5
4	4-FC <sub>6</sub> H <sub>4</sub>	H	<b>1e</b>	71%	91 : 9
5	4-ClC <sub>6</sub> H <sub>4</sub>	H	<b>1f</b>	68%	90 : 10
6	4-BrC <sub>6</sub> H <sub>4</sub>	H	<b>1g</b>	68%	91 : 9
7	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	H	<b>1h</b>	20%	92 : 8
8	Ph	CH <sub>3</sub>	<b>1i</b>	70%	95 : 5
9	Ph	<i>n</i> -Bu	<b>1j</b>	72%	91 : 9
10 <sup>c</sup>	Ph	Ph	<b>1k</b>	56%	99 : 1

<sup>a</sup> Reaction conditions: Alkynes (0.48 mmol) and BrCN (0.40 mmol) in ClCH<sub>2</sub>CH<sub>2</sub>Cl (1.6 mL) were heated in the presence of GaCl<sub>3</sub> (10 mol%). <sup>b</sup> Determined by NMR. <sup>c</sup> Reaction carried out in toluene at 100 °C.

The alkynophilicity<sup>18</sup> of trivalent GaX<sub>3</sub> leading to the formation of cationic vinylgallium species is well known, and some notable synthetic applications have been demonstrated.<sup>19</sup> To gain insight into the present bromocyanation of alkynes, we carried out an NMR study on a stoichiometric reaction. When GaCl<sub>3</sub> was added to a solution of BrCN in CDCl<sub>3</sub> at room temperature, the signal of CN (δ 76.1 ppm) in BrCN shifted to a new peak at δ 88.2 ppm. The downfield shift of the CN peak suggested the possibility of the formation of a complex between BrCN and GaCl<sub>3</sub>.<sup>20</sup> When an equimolar amount of 1-phenyl-1-hexyne was added to a CDCl<sub>3</sub> solution of this complex at room temperature, the quantitative formation of the bromocyanation product **1j** (Z:E = 98:2) coordinated with GaCl<sub>3</sub> was observed, with the signal of the CN moiety being observed at δ 149.5 ppm. This result clearly shows that electrophilic addition of the BrCN

and GaCl<sub>3</sub> complexes to alkynes<sup>21</sup> occurs, even at room temperature, and a high temperature is required in the catalytic reaction conditions to release GaCl<sub>3</sub> from cyano moiety of the adduct.

The synthetic utility of (*Z*)-β-bromo-α,β-unsaturated nitriles obtained from the bromocyanation of alkynes was demonstrated by the cross-coupling reactions of the representative product **1a** (Scheme 2). The Stille coupling reactions of **1a** with organostannanes afforded the stereo-defined structures **2** or **3** in good yields. The Sonogashira coupling reaction of **1a** with phenylacetylene gave enyne **4** quantitatively, with complete stereoselectivity. The nickel-catalysed reductive homo-coupling of **1a** produced 3,4-diphenyl-2,4-hexadiene-1,6-dinitrile **5**.<sup>22</sup> Moreover, we demonstrated the synthetic utility of **1a** and its derivative **3** in the preparation of the biologically active heterocycles **6**<sup>23</sup> and **7**.<sup>11</sup>



(a) (4-CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>)SnBu<sub>3</sub>, Pd(PPh<sub>3</sub>)<sub>4</sub>, CuI, dioxane, 100 °C, 8h. (b) BzSnBu<sub>3</sub>, Pd(OAc)<sub>2</sub>, PPh<sub>3</sub>, dioxane, 100 °C, 12 h. (c) Phenylacetylene, Pd(PPh<sub>3</sub>)<sub>4</sub>, CuI, Et<sub>3</sub>N, THF, rt, 5 h. (d) NiBr<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>, PPh<sub>3</sub>, Zn, dioxane, 80 °C, 6 h. (e) Ethyl thioglycolate, NaOEt, EtOH, 70 °C, 12 h. (f) 1,3-Dimethoxybenzene, Cu(OTf)<sub>2</sub>, ClCH<sub>2</sub>CH<sub>2</sub>Cl, H<sub>2</sub>O, 80 °C, 15 h. Ar = 2,4-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>.

**Scheme 2** Transformation of **1a**.

In summary, we developed gallium(III)-catalysed bromocyanation of alkynes using cyanogen bromide. This method enables the regio- and stereoselective introduction of the synthetically useful Br and cyano functionalities to carbon-carbon triple bonds in single operation. Further investigations into the reaction mechanism, substrate scope, and the synthetic application are currently underway in our laboratory.

**Acknowledgment.** This work is financially supported by a Grant-in-Aid for Scientific Research from MEXT. M. M. thanks the JSPS Research Fellowships for Young Scientists.

## Notes and references

- 1 B. M. Trost, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 259.
- 2 (a) H.-J. Arpe, in *Industrial Organic Chemistry*, 5th ed.; Wiley-VCH: Weinheim, **2010**, p 312; (b) A. R. Katritzky; O. Meth-Cohn; C. W. Rees, in *Comprehensive Organic Functional Group Transformations*; Pergamon, Oxford, **1995**, Vol. 3, p 614; (c) B. Cornils; W. A. Herrmann, in *Applied Homogeneous Catalysis with Organometallic*

- Compounds*; VCH, Weinheim, **1996**, Vol. I, p 476; (d) M. Beller; C. Bolm, in *Transition Metals for Organic Synthesis 2nd ed.*; Wiley-VCH, Weinheim, **2004**, p 151.
- 3 For selected examples, see: (a) K. Nozaki; N. Sato; H. Takaya, *J. Org. Chem.* **1994**, *59*, 2679; (b) Y. Nakao; S. Oda; T. Hiyama, *J. Am. Chem. Soc.* **2004**, *126*, 13904; (c) Y. Nakao; K. S. Kanyiva; S. Oda; T. Hiyama, *J. Am. Chem. Soc.* **2006**, *128*, 8146; (d) Y. Kobayashi; H. Kamisaki; R. Yanada; Y. Takemoto, *Org. Lett.* **2006**, *8*, 2711; (e) Y. Nakao; A. Yada; S. Ebata; T. Hiyama, *J. Am. Chem. Soc.* **2007**, *129*, 2428; (f) Y. Hirata; T. Yukawa; N. Kashiwara; Y. Nakao; T. Hiyama, *J. Am. Chem. Soc.* **2009**, *131*, 10964; For carbocyanation to alkenes or 1,2-dienes, see: (g) Y. Nishihara; Y. Inoue; M. Itazaki; K. Takagi, *Org. Lett.* **2005**, *7*, 2639; (h) Y. Nakao; Y. Hirata; T. Hiyama, *J. Am. Chem. Soc.* **2006**, *128*, 7420; (i) Y. Nakao; S. Ebata; A. Yada; T. Hiyama; M. Ikawa; S. Ogoshi, *J. Am. Chem. Soc.* **2008**, *130*, 12874; (j) Y. Yasui; H. Kamisaki; Y. Takemoto, *Org. Lett.* **2008**, *10*, 3303; (k) M. P. Watson; E. N. Jacobsen, *J. Am. Chem. Soc.* **2008**, *130*, 12594; (l) Y. Hirata; T. Inui; Y. Nakao; T. Hiyama, *J. Am. Chem. Soc.* **2009**, *131*, 6624.
- 4 (a) N. Chatani; T. Hanafusa, *J. Chem. Soc., Chem. Commun.* **1985**, 838; (b) N. Chatani; T. Takeyasu; N. Horiuchi; T. Hanafusa, *J. Org. Chem.* **1988**, *53*, 3539; For 1,2-dicyanation, see: (c) S. Arai; T. Sato; Y. Koike; M. Hayashi; A. Nishida, *Angew. Chem. Int. Ed.* **2009**, *48*, 4528.
- 5 N. Chatani; N. Horiuchi; T. Hanafusa, *J. Org. Chem.* **1990**, *55*, 3393.
- 6 Y. Obora; A. S. Baleta; M. Tokunaga; Y. Tsuji, *J. Organomet. Chem.* **2002**, *660*, 173.
- 7 (a) M. Sugimoto; A. Yamamoto; M. Murakami, *J. Am. Chem. Soc.* **2003**, *125*, 6358; (b) M. Sugimoto; A. Yamamoto; M. Murakami, *Angew. Chem. Int. Ed.* **2005**, *44*, 2380.
- 8 (a) I. Kamiya; J. Kawakami; S. Yano; A. Nomoto; A. Ogawa, *Organometallics* **2006**, *25*, 3562; (b) Z. Zhang; L. S. Liebeskind, *Org. Lett.* **2006**, *8*, 4331; (c) Y. T. Lee; S. Y. Choi; Y. K. Chung, *Tetrahedron Lett.* **2007**, *48*, 5673.
- 9 For uncatalysed bromocyanation of ynamines, see: (a) N. V. Lukashev; A. V. Kazantsev; A. A. Borisenko; I. P. Beletskaya, *Tetrahedron* **2001**, *57*, 10309. For copper-promoted iodocyanation of (perfluoroalkyl)-alkynes, see: (b) P. Moreau; A. Commeyras, *J. Chem. Soc., Chem. Commun.* **1985**, 817.
- 10 For bromocyanation of enamines, see: (a) F. Raffaello; R. Silvano; B. Giuseppe, *Gazz. Chim. Ital.* **1961**, *91*, 841; (b) W. Verboom; G. W. Visser; D. N. Reinhoudt, *Tetrahedron* **1982**, *38*, 1831; (c) N. D. Kimpe; R. Verhé; L. D. Buyck; N. Schamp, *Chem. Ber.* **1983**, *116*, 3846.
- 11 For synthetic applications of  $\beta$ -acyl- $\alpha,\beta$ -unsaturated nitriles, see: (a) M. Murai; S. Kawai; K. Miki; K. Ohe, *J. Organomet. Chem.* **2007**, *692*, 579; (b) M. Murai; K. Miki; K. Ohe, *J. Org. Chem.* **2008**, *73*, 9174.
- 12 Cyanogen bromide (mp 52 °C) is a commercially available reagent and easy to handle. However, the handling of cyanogen bromide requires special care to avoid inhalation of hydrogen cyanide, and all operations should be carried out in a well-fumed hood.
- 13 (a) C. C. Price; J. A. Pappalardo, *J. Am. Chem. Soc.* **1950**, *72*, 2613; (b) H. Martens; F. Janssens; G. Hoornaert, *Tetrahedron* **1975**, *31*, 177; (c) H. Zhou; C. Zeng; L. Ren; W. Liao; X. Huang, *Synlett* **2006**, 3504; For electrophilic cyanation of arenes with BrCN, see: (d) P. H. Gore; F. S. Kamounah; A. Y. Miri, *Tetrahedron* **1979**, *35*, 2927.
- 14 The cyano group was selectively introduced at the terminal position, no regioisomers of **1a** being detected.
- 15 The use of polar solvents, such as MeCN, dioxane, THF, and DMF resulted in complete recovery of phenylacetylene.
- 16 The structure of the major stereoisomer of **1e** was unambiguously determined by X-ray crystallography. See electronic supplementary information for the detail of X-ray crystal analysis data. CCDC-796829 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- 17 Determination of regio- and stereoselectivity of the major adduct was based on derivatization to a stereo-defined compound by reduction of **1h** with DIBAL-H and NaBH<sub>4</sub> (See electronic supplementary information).
- 18 For selected reviews, see: (a) A. Fürstner; P. W. Davies, *Angew. Chem. Int. Ed.* **2007**, *46*, 3410. (b) K. Ohe; K. Miki, *J. Synth. Org. Chem., Jpn.* **2009**, *67*, 1161 and references therein.
- 19 For a review in this area, see: R. Amemiya; M. Yamaguchi, *Eur. J. Org. Chem.* **2005**, 5145.
- 20 Although precise species are not clear at present, the formation of  $\delta^{\text{Br}}\text{--}\delta^{\text{CN}}\text{CN-GaCl}_3$  or  $[\text{CN}]^+[\text{GaCl}_3\text{Br}]^-$  is most likely.
- 21 We suppose that electrophilic addition of a positively charged CN followed by intramolecular nucleophilic attack of bromide in the intimate complex to the resulting vinyl cation center in syn fashion provides a Z-adduct. Precise mechanism awaits further investigation.
- 22 For application of **5** to electrophotographic photoreceptor, see: K. Watanabe, *Jpn. Kokai Tokkyo Koho* **1997**, JP 09006027 [*Chem. Abstr.* **1997**, *126*, 205452].
- 23 For selected examples, see: (a) A. V. Bogolubsky; S. V. Ryabukhin; A. S. Plaskon; S. Stetsenko; D. M. Volochnyuk; A. A. Tolmachev, *J. Comb. Chem.* **2008**, *10*, 858. (b) Z. Brzozowski; J. Sławiński; F. Saçzewski; T. Sanchez; N. Neamati, *Eur. J. Med. Chem.* **2008**, *43*, 1188.